
Organic matter composition and thermal maturity assessment by the use of the Palynomorph Darkness Index method: a case study from the middle-upper Cenomanian OAE Black-Shales (Göynük-Sünnet section - NW Turkey).

Simone Bonciani^{*1}, Enrico Capezzuoli², Ismail Omer Yilmaz³, Nicoletta Buratti⁴, and Amalia Spina^{†5}

¹Department of Earth Sciences, University of Florence – Italy

²Department of Earth Sciences, University of Florence – Italy

³Middle East Technical University, Ankara – Turkey

⁴TotalEnergies SE, Pau Cedex – TotalEnergies S.E. – France

⁵Department of Physics and Geology, University of Perugia – Italy

Abstract

A palynological study from the Upper Cretaceous Uzumlu Member of Yenipazar Formation (NW Turkey) of the Mudurnu-Goynuk Basin was yielded with the aim to assess the organic matter composition and to evaluate the thermal maturity. For this purpose, in the pelagic succession of the Göynük-Sünnet section (north-western Turkey) located in a rift basin along the Sakarya continental margin twenty-seven black shale levels of middle-upper Cenomanian age were sampled. Such levels intercalated with marly shale/calcareous marl rhythmic alternations and chronostratigraphically constrained to Cenomanian stage by the planktonic foraminifer *Rotalipora cushmani*. Additionally, the planktonic foraminifer *Wheitenella archeocretacea* was recorded across the Cenomanian-Turonian boundary (Yilmaz, 2008) allowing to associate these black shales to mid-Cenomanian Oceanic Anoxic Event and to the Bonarelli Event OAE2. The organic matter content analysed was mainly of continental origin showing abundance of oxidised and partially oxidised ligneous particles, pollen, spores and less frequent marine elements as dinoflagellates and acritarchs. To evaluate the thermal maturity of organic matter, optical semi-quantitative method as the Palynomorph Darkness Index (PDI; *Goodhue and Clayton, 2010*) was used. PDI was measured on smooth, unfolded trilete spores and bisaccate pollens and calibrated with other thermal index as the Spore Colour Index (SCI; Waples, 1985; Marshall, 1991). PDI value varies based on the taxon measures. Generally, trilete spore shows PDI values ranging from 13% to 34%. The SCI observations (SCI 3.2 to 4.2) also indicate that organic matter content experienced a very immature stage with conversion vitrinite reflectance values (Clayton, et al., 2017) below 0.40%. In this study for the first time Palynomorph Darkness Index has been calibrated on Upper Cretaceous deposits thus opening new scenarios to other types of studies as sedimentary basins thermal history assessment. Moreover, further studies are in

^{*}Corresponding author: simone.bonciani1@stud.unifi.it

[†]Speaker

progress to rigorously calibrate the microflora with foraminifera biozones as well as with the OAEs.

Clayton, G., Goodhue, R., Abdelbagi, S. T. & Vecoli, M., 2017. *Correlation of Palynomorph Darkness Index and Vitrinite reflectance in a submature Carboniferous well section in northern Saudi Arabia*. *Revue de micropaléontologie*, Issue 60, pp. 411-416.

Goodhue, R., & Clayton, G., 2010. *Palynomorph Darkness Index (PDI) – a new technique for assessing thermal maturity*. *Palynology*, 34(2), pp. 147-156.

Marshall, J. E. A., 1991. *Quantitative spore colour*. *Journal of the Geological Society of London*, Volume 148, pp. 223-233

Waples, D., 1985. *Geochemistry in petroleum Exploration*: Boston, IHRDC, 232 p.

Yilmaz, I., Ö., 2008. *Cretaceous Pelagic Red Beds and Black Shales (Aptian-Santonian), NW Turkey: Global Oceanic Anoxic and Oxidic Events*. *Turkish Journal of earth Sciences*, pp. 263-296.

Keywords: Thermal maturity of Organic Matter, Palynomorph Darkness Index, Mid, Cenomanian, Turkey